



# An Interesting case of PUO



# About the patient

61/F admitted for evaluation of fever since March 2008:

- Fever high grade intermittent followed by sweating  
no evening rise or specific pattern  
no chills or rigors  
body pain+  
arthralgia involving large joints without swelling ,small joint involvement, hand joint involvement or early morning stiffness  
loss of appetite+ no significant weight loss



She was treated in her native place with only temporary relief

- Breathlessness insidious onset- exertional, gradually worsened to Class IV with orthopnoea ; no PND
- Cough with small quantity mucoid sputum, no hemoptysis
- Negative past history except for a suppurative left axillary adenitis which resolved with treatment
- Negative family history; three healthy siblings

She was hospitalized and the following investigations were done:

# Investigations:2/4/08

Hb 12.6gm/dl

TC 9000 cells/cu mm

ESR 22/45

Urine: SG 1.005

pH 7.0

WBC 1+

RBC nil

Nitrite neg

Protein neg

Glucose neg

Ketones neg

Urobil, neg

ECG: Sinus tachycardia

CXR cardiomegaly; lung fields clear

Widal neg

US Abdomen: Hepatomegaly

RK 10.8x4

LK 10.7x4

No ascites

HIV I&II neg

RFT&LFT normal

HBsAg neg

Blood C/S no growth



# Contd..

Echo: Pericardial effusion 500ml with early tamponade; valves, chambers normal; no SWMA; N LV function

TFT: T3 71.69ng/dl(N 80-180)

T4 3.6mgm/dl(N 4.5-11.5)

TSH 100.960mIu/ml(N 0.35-5.5)

Cardiologist suggested medical management of effusion.

Patient was started on ATT on 4/4/08 along with Thyroxine 100mcg and discharged with a provisional diagnosis of Tuberculous Pericardial effusion, and Hypothyroidism.

CT Abdomen normal

However, she discontinued ATT and got admitted on 4/5/08.

# On admission

O/E Obese lady

- Febrile
- Dyspnoeic and tachypnoeic
- No cyanosis, clubbing
- No pallor, adenopathy
- Oral ulcers+
- No skin, hair, nail or eye changes; no bony tenderness or joint swelling/deformities
- JVP not elevated; no pedal edema or facial puffiness
- Tachycardic, BP 120/70, all peripheral pulses+
- CVS: Heart sounds were normal; no gallop or murmurs
- RS: Trachea in midline; NVBS; Coarse crepitations in Rt interscapular, infrascapular, axillary, infra-axillary areas with diffuse rhonchi
- Abd : No ascites or organomegaly
- CNS:N

# Investigations done :4/5/08

Hb 8.1gm/dl

TC 8900 cells/cu mm

Widal negative

ESR 20/42

RFT & LFT N

Platelets 3.15 lakh

MCV 77.6

MCHC 33.6

MCH 26.1

Peripheral smear: microcytic hypochromic anemia

Urine analysis: albumin-nil

pus cells-3-4/hpf

RBC-nil

bacteria-nil


ECG Sinus Tachycardia T inv in V2-4


Echo Pericardial effusion; no tamponade


Cardiologist opinion conservative management

CXR Cardiac silhouette enlarged with bilateral patchy infiltrates more in right lower zone with obliteration of costo and cardiophrenic angles











# Contd..6/5/08

Sputum 3 samples negative for AFB

HIV 1&2 negative

Sputum C/S Staph aureus

ASO +ve 400 IU/ml

CRP +ve 96mg/dl

ANA +ve


CT Chest: B/L Pleural effusion


homogenous airspace opacity-posterior  
segment of right UL

and superior segment of right lower lobe











# Contd..

Provisional Diagnosis:

SLE

- \*Serositis- Bilateral pleural& pericardial effusions

- \*Rt lower lobe consolidation:

- ?Infective

- ?Acute Lupus Pneumonitis



# Treatment

- Rheumatologist opinion:

SLE with patchy pneumonitis and serositis

?Infective

?Lupus pneumonitis

For Antibiotics and Prednisolone

Repeat Sputum C/S on 14.5.08 grew Staph aureus sensitive to Vancomycin & Vancomycin started on 18.5.08

Patient continued to be febrile and tachypnoeic

Developed elevated renal parameters on 5<sup>th</sup> day and vancomycin stopped

Dyspnoea worsened and was shifted to ICU for respiratory support on 21.5.08



# RFT

	8.5.08	22.5	23.5 renal lab	24.5	27.5	29.5
Urea	25	82	121	55	140	132
Creat	0.6	2.4	3.1	2.6	2.1	1.5



# ABG

pH 7.495

pCO<sub>2</sub> 20.6

pO<sub>2</sub> 67.4

HCO<sub>3</sub> 15.5

BE(ecf) -7.8

O<sub>2</sub> sat 95.2%

CO<sub>2</sub> 16.1mmol/L

Na 135 meq/L

K 4.3 meq/L

# At ICU

- Anti ds DNA 30.4 U/ml

(neg <20 U/ml

pos >20 U/ml)

- Pericardiocentesis was done:

Sugar 79mg/dl

Protein 4.4gm/dl

Cells RBC-120 cells; Lymphocytes-8 cells

C/S no growth

Smear neg for AFB

ADA (fluid) 40.4 U/L

Serum ADA 69.9 U/L

- Urine C/S grew Pseudomonas

Patient was treated with antibiotics, low dose steroids (oral pred 10mg/d) and fluid management

Her metabolic parameters improved; did not require ventilation and shifted back to ward



# Contd..

Patient continued to be febrile and tachypnoeic despite antibiotics

Minimal sputum production; dry cough

Repeated sputum C/S and AFB neg

Rheumatologist reviewed and suggested parenteral steroids

Started on IV Methyl Pred 1gm/d x 5 days

Patient improved from second dose.



# Final Diagnosis

LUPUS PNEUMONITIS



# Pleuro pulmonary manifestations of SLE

- Lupus pneumonitis
- Lymphocytic interstitial pneumonitis
- Pulmonary hemorrhage
- Pulmonary embolism associated with aPL
- Pulmonary hypertension
- Pleuritis
- Weakness of diaphragm

# Lupus pneumonitis

- Acute:

- 12% of active lupus

- fever, pleuritic pain, dyspnoea, cough, cyanosis

- B/L pulmonary infiltrates and effusion

- HP: alveolar damage, interstitial edema, hyaline membranes

- perivascular lymphocytic & plasma cell infiltrates- clear or persist causing PFT abn

- Chronic:

- Similar to other interstitial lung diseases

- Cough-nonproductive, dyspnoea, basilar rales and abn PFT with persistent infiltrates

- HP: fibrosis, necrosis, plasma cell infiltration with

- IF: immune complex in alveolar wall



# Lupus pneumonitis contd..

Diagnosis is one of exclusion

- D/D

  - Infective consolidation

  - Pulmonary hemorrhage

- Treatment:

  - acute- steroids, immunosuppressants if steroid unresponsive

  - chronic- asymptomatic: no treatment; poor prognosis if PFT abn

- Prognosis:

  - poor; 50% mortality

  - sequelae for survivors is severe restrictive lung disease



THANK YOU